



---

---

Exercises and Complements for the Introduction to Physics II  
for Students  
of Biology, Pharmacy and Geoscience

---

---

Sheet 10 / 06.05.2021

**Solutions**

**Exercise 37.**

For the angular magnification of the magnifying glass we have:

$$\Gamma = \frac{s_0}{f} = \frac{0.25 \text{ m}}{\frac{1}{12 \text{ dpt}}} = 3$$

**Exercise 38.**

The following applies to the intensity of the light  $I_2$  that passes through the polarization film:

$$I_2 = I_1 \cos^2 \theta$$

Here  $I_1$  is the intensity of the light before it hits the film and  $\theta$  is the angle that the transmission axis forms with the horizontal. This results in:

$$\arccos \sqrt{\frac{I_2}{I_1}} = \arccos \sqrt{0.15} = 67.2^\circ$$

**Exercise 39.**

- (a) true, because the polarization axes of the last two filters are rotated by  $90^\circ$  to each other
- (b) true, because after passing a polarizing filter twisted by  $45^\circ$  only exactly half of the intensity comes through ( $\cos^2 45^\circ = 0.5$ )
- (c) false, because the polarization axes of the filters are shifted by  $90^\circ$ , this is impossible
- (d) true, since the light is previously unpolarized
- (e) true, because the polarization axes of the filters are always shifted by  $45^\circ$  and the light is unpolarized in the beginning

**Exercise 40.**

(a) See script 507-3. angle of incidence  $\alpha_B$  for full polarization from Brewster's law:

$$\tan \alpha_B = \frac{n_{\text{glass}}}{n_{\text{air}}} \quad \Rightarrow \quad \alpha_B = 55.41^\circ$$

(b) See script 507-7.

$$\alpha = \varphi \cdot c \cdot d \quad \Rightarrow \quad c = \frac{\alpha}{\varphi \cdot d} = 3.0 \text{ g/l}$$

(c)  $\alpha = 0$ , since the same number of levorotatory and dextrorotatory molecules are present.